

Real Commuting Differential Operators Connected with Two-Dimensional Abelian Varieties

A.E. Mironov

1 Introduction

In [1] A. Nakayashiki constructed commutative rings of $(g! \times g!)$ -matrix partial differential operators in g variables (see also [2]). The common vector eigenfunctions and eigenvalues of these operators are parametrized by points of some principally polarized abelian variety of dimension g with a nonsingular theta-divisor. Each operator corresponds to some meromorphic function (spectral function) on this abelian variety with a pole on the theta-divisor. Henceforth these operators are referred to as *Nakayashiki operators*.

Some explicit formulas for Nakayashiki operators with $g = 2$ were obtained in [3]. Using these formulas, we find smooth real operators.

Theorems 1 and 2 are the main results of this article.

Theorem 1. *For $g = 2$, there are no Nakayashiki operators with smooth real doubly periodic coefficients but there are Nakayashiki operators with real singular doubly periodic coefficients.*

This theorem is an analog of the theorem of Feldman, Knorrer, and Trubowitz [4] who demonstrated that a two-dimensional Schrödinger operator without magnetic field which has a smooth doubly periodic real potential can be finite-gap only at one energy level; i.e., the Bloch functions (the eigenfunctions of the Schrödinger operator and the translation operators by periods) may be parametrized by a Riemann surface of finite genus only at one energy level. Theorem 1 means that there is no smooth real Nakayashiki operators that are finite-gap at all energy levels. Nevertheless, there exist real Nakayashiki operators with singular coefficients which are finite-gap at every energy level.

We take as the abelian variety the Jacobi variety of a Riemann surface of genus 2 with real branching points. In this case the symmetric matrix Ω of the periods of basis abelian differentials has purely imaginary components [5]. Introduce the magnetic translation operators T_1^* and T_2^* :

$$T_1^* \varphi(y) = \varphi(y + e_1) \exp(2\pi y_1), \quad T_2^* \varphi(y) = \varphi(y + e_2) \exp(2\pi y_2),$$

where $y = (y_1, y_2)$, e_j is the j th row of the imaginary part of the period matrix Ω . The magnetic translation operators differ from the translation operators only by an exponential twist. The arguments of exponential functions in the magnetic translation operators are chosen so that $A_i(y+e_j) - A_i(y) = 2\pi\delta_{ij}$, where (A_1, A_2) is a vector-potential of the magnetic field [6]. Then the operators T_j^* commute with the covariant differentiation operators $\partial_{y_i} - A_i$. The operators T_1^* and T_2^* commute with one another. This is a consequence of the fact that in our case the magnetic flux through the elementary cell formed by the vectors e_1 and e_2 vanishes. In the general case we have $T_1^*T_2^* = T_2^*T_1^* \exp(ie\Phi)$, with e the charge and Φ the magnetic flux [6], and the operators T_1^* and T_2^* commute if $\frac{e\Phi}{2\pi}$ is integral.

An eigenfunction of a matrix differential operator is called a *magneto-Bloch function* if its components are eigenfunctions of the magnetic translation operators. Denote the theta-function of the abelian variety $\mathbb{C}^2/\{\mathbb{Z}^2 + \Omega\mathbb{Z}^2\}$ by $\theta(z)$, with $z = (z_1, z_2)$.

Theorem 2. *There exist Nakayashiki operators with smooth real coefficients. The diagonal of the operator H corresponding to the function $\partial_{z_1}^2 \ln \theta(z) + \partial_{z_2}^2 \ln \theta(z)$ is composed of Schrödinger operators of the form*

$$H_{11} = (\partial_{y_1} - A_1)^2 + (\partial_{y_2} - A_2)^2 + u(y), \quad H_{22} = (\partial_{y_1} - \tilde{A}_1)^2 + (\partial_{y_2} - \tilde{A}_2)^2 + \tilde{u}(y)$$

with doubly periodic magnetic fields $\text{rot}(A_1, A_2, 0)$ and $\text{rot}(\tilde{A}_1, \tilde{A}_2, 0)$ and with doubly periodic potentials

$$u(y + e_j) = u(y), \quad \tilde{u}(y + e_j) = \tilde{u}(y).$$

The components of the vector-potentials satisfy the equalities

$$A_i(y + e_j) - A_i(y) = \tilde{A}_i(y + e_j) - \tilde{A}_i(y) = 2\pi\delta_{ij}.$$

The magneto-Bloch functions of H at each energy level are parametrized by Riemann surfaces of finite genus. The components of H commute with T_1^* and T_2^* .

We also indicate Nakayashiki operators of simplest form. For example, the operators L and L_1 corresponding to the functions $\partial_{z_1}^2 \log \theta(z)$ and $\partial_{z_1} \partial_{z_2} \log \theta(z)$ are as follows:

Lemma 1. *The following hold:*

$$L = \begin{pmatrix} -\partial_{x_1}^2 + c_1 \partial_{x_2} + U & W \\ \frac{V}{c_2}(-\partial_{x_1}^2 + c_1 \partial_{x_2} + U - c_3) & -\partial_{x_1}^2 - c_1 \partial_{x_2} + \tilde{U} + \frac{WV}{c_2} \end{pmatrix},$$

where

$$U = \partial_{x_1}^2 \ln V + (\partial_{x_1} \ln V + c_4)^2 - c_1(\partial_{x_2} \ln V + c_5) + c_3,$$

$$\tilde{U} = \partial_{x_1}^2 \ln W + (\partial_{x_1} \ln W - c_4)^2 + c_1(\partial_{x_2} \ln W - c_5) + c_3;$$

and

$$L_1 = \begin{pmatrix} -\partial_{x_1} \partial_{x_2} + U_2 \partial_{x_1} + c_6 \partial_{x_2} + U_1 & W_1 \\ \frac{V_1}{c_2} (-\partial_{x_1}^2 + c_1 \partial_{x_2} + U - c_3) & -\partial_{x_1} \partial_{x_2} + \tilde{U}_2 \partial_{x_1} - c_6 \partial_{x_2} + \tilde{U}_1 + \frac{WV_1}{c_2} \end{pmatrix},$$

where

$$U_1 = \partial_{x_1} \partial_{x_2} \ln V + (\partial_{x_1} \ln V + c_4)(\partial_{x_2} \ln V + c_5) - U_2(\partial_{x_1} \ln V + c_4) - c_6 \partial_{x_2} \ln V + c_7,$$

$$\tilde{U}_1 = \partial_{x_1} \partial_{x_2} \ln W + (\partial_{x_1} \ln W - c_4)(\partial_{x_2} \ln W - c_5) - \tilde{U}_2(\partial_{x_1} \ln W - c_4) + c_6 \partial_{x_2} \ln W + c_7,$$

$$W_1 = \frac{c_6}{c_1} W - \frac{1}{2c_1} \partial_{x_1} W, \quad V_1 = \frac{c_6}{c_1} V + \frac{1}{2c_1} \partial_{x_1} V,$$

$$U_2 = \frac{1}{2c_1} (U + c_8), \quad \tilde{U}_2 = -\frac{1}{2c_1} (\tilde{U} + c_8),$$

and c_j are some constants (see (15)–(18)).

Observe that the coefficients of ∂_{x_2} in the 11- and 22-components of these operators are constants and the coefficients of ∂_{x_1} in L vanish. Moreover, all coefficients of L and L_1 are rationally expressed in terms of V and W and their derivatives. There are no similar relations between the coefficients of the operators in [3].

We indicate partial solutions to the system of nonlinear equations $[L, L_1] = 0$ in V and W . The solutions are given by (8) and (9).

Theorem 3. *The coefficients of the Nakayashiki operators are rationally expressible in terms of the coefficients V and W of L and their derivatives.*

Observe that the 11-components of the operators commute modulo a heat operator (Lemma 8); i.e., for arbitrary two 11-components A and B there is an operator C such that

$$[A, B] = C(-\partial_{x_1}^2 + c_1 \partial_{x_2} + U - c_3).$$

The coefficients of the Nakayashiki operators cannot satisfy evolution equations like the Kadomtsev–Petviashvili hierarchy (KP)

$$[\partial_{t_n} - L_n, \partial_{t_m} - L_m] = 0.$$

Indeed, to the space variables there correspond g linearly independent rectilinear windings on the g -dimensional torus. To the time variables there also correspond rectilinear windings on the torus; moreover, they are linear combinations of spatial windings, since the dimension of the torus coincides with the number of the space variables. Consequently, the time derivatives of the coefficients of operators can be expressed linearly in terms of the derivatives with respect to the space variables and, changing variables, we can reduce “evolution” equations to commutation equations for Nakayashiki operators. In the particular case of $g = 1$ the finite-gap solutions of the KP hierarchy (the so-called stationary solutions) are not interesting, since replacement of differentiation with respect to time by differentiation with respect to the space variables takes the equations of the hierarchy into the commutation equations $[L_n, L_m] = 0$ for the operators.

Let us show that Nakayashiki’s construction [1] does not lead to evolution equations. Below all notations until Section 2 are taken from [1]. Introduce two functions

$$\begin{aligned}\varphi_1 &= \frac{\theta(z + (c' - x'd' - x_0, x'))}{\theta(z)} \exp, \\ \varphi_2 &= \frac{\theta(z + (c' - x'd' - x_0, x') + c'')\theta(z - c'')}{\theta^2(z)} \exp, \quad c'' \in \mathbb{C}^2, c'' \notin \mathbb{Z}^2 + \Omega\mathbb{Z}^2, \\ \exp &= \exp\left(\sum_{i=0}^1 \sum_{n \geq \delta_{i0}} t_{n,(i)} \frac{(-1)^n}{n!} (u_{n,(i)}(z) + d_i(1 - \delta_{i0})u_{n+1,(0)}(z))\right),\end{aligned}$$

where x' and x_0 are the space variables, $u_{n,(i)}$ are the derivatives of the logarithm of the theta-function, c' , d_i , and $\delta_{i,0}$ are constants, $x_0 = t_{1,(0)}$, and $x' = t_{0,(1)}$. These functions determine a basis for the free module B_{ct} over $\mathcal{D}_t$, where $\mathcal{D}_t$ is the ring of differential operators in the variables x_0 and x' with analytic coefficients depending on $t_{n,(i)}$ in a neighborhood of 0. In [1] some embedding ι was constructed of the $\mathcal{D}_t$ -module B_{ct} into the ring of pseudodifferential operators. Equations (6.8) of [1] read:

$$\frac{\partial W_i}{\partial t_\beta} + W_i \partial^\beta = \sum_{j=1}^2 B_{i,\beta,j} W_j,$$

where $W_i = \iota(\varphi_i)$ and $B_{i,\beta,j} \in \mathcal{D}_t$. Observe that the image of $\varphi \in B_{ct}$ under ι depends only on $\frac{\varphi}{\exp}$. Consequently, the operators W_i are independent of time, and $\frac{\partial W_i}{\partial t_\beta} = 0$. Hence, (6.8) of [1] are not evolution equations.

In Section 2 we recall Nakayashiki's construction and prove Theorems 1 and 2.

In Section 3 we prove Lemma 1 and Theorem 3.

The author is grateful to I. A. Taĭmanov for posing the problem as well as for useful discussions and remarks.

2 Smooth Real Operators

We start with recalling Nakayashiki's construction of the Baker–Akhiezer module M_c . Then we prove Theorem 1. Next we introduce a hyperelliptic surface Γ of genus 2 with real branching points and take a canonical basis of cycles on it by utilizing the scheme of [5]. Thereafter we introduce four real two-dimensional tori T_j , $1 \leq j \leq 4$, in X (the Jacobi variety of Γ). The theta-function takes real values on these tori. In Lemma 3 we prove that the theta-function has no zeros on T_1 . In Lemma 4 we find the intersection points of the theta-divisor and the translated theta-divisor. The proof of Theorem 2 consists in verifying that the coefficients of the Nakayashiki operators of the second and third orders from Propositions 1–3 of [3], with changes made below, are real and smooth.

Consider the principally polarized complex abelian variety $X = \mathbb{C}^2 / \{\mathbb{Z}^2 + \Omega\mathbb{Z}^2\}$, where Ω is a symmetric 2×2 -matrix with $\text{Im}\Omega > 0$. The theta-function is defined by the series

$$\theta(z) = \sum_{n \in \mathbb{Z}^2} \exp(\pi i \langle \Omega n, n \rangle + 2\pi i \langle n, z \rangle), \quad z \in \mathbb{C}^2,$$

where $\langle n, z \rangle = n_1 z_1 + n_2 z_2$. It has the periodicity properties

$$\theta(z + \Omega m + n) = \exp(-\pi i \langle \Omega m, m \rangle - 2\pi i \langle m, z \rangle) \theta(z), \quad m, n \in \mathbb{Z}^2.$$

Let $\mathcal{D}$ be the ring $\mathcal{O}[\partial_{x_1}, \partial_{x_2}]$ of differential operators, where $\mathcal{O}$ is the ring of analytic functions (of x) in a neighborhood of $0 \in \mathbb{C}^2$. In [1] Nakayashiki introduced the Baker–Akhiezer module M_c over $\mathcal{D}$ which consists of functions of the form

$$f(z, x) \exp(-x_1 \partial_{z_1} \log \theta(z) - x_2 \partial_{z_2} \log \theta(z)).$$

The function $f(z, x)$ is meromorphic on $\mathbb{C}^2 \times U_f$, where U_f is some neighborhood of $0 \in \mathbb{C}^2$, has a pole in the theta-divisor Θ (the zeros of the theta-function $\theta(z)$), and possesses the periodicity property

$$f(z + \Omega m + n, x) = \exp(-2\pi i \langle m, c + x \rangle) f(z, x), \quad m, n \in \mathbb{Z}^2, \quad c = (c_1, c_2) \in \mathbb{C}^2.$$

The following was proven in [1]:

Nakayashiki's Theorem. *If Θ is a nonsingular variety and $c \neq 0$ then M_c is a free $\mathcal{D}$ -module of rank 2.*

Fix a basis $\Phi_c = (\phi_{1c}(z, x), \phi_{2c}(z, x))^\top$ for the $\mathcal{D}$ -module M_c . Denote by $\mathcal{A}_\Theta$ the ring of meromorphic functions on X with a pole in Θ . Take $\lambda(z) \in \mathcal{A}_\Theta$. Since M_c is a free $\mathcal{D}$ -module, there is a unique (2×2) -matrix operator $L_{\Phi_c}(\lambda)$ with components in $\mathcal{D}$ such that

$$L_{\Phi_c}(\lambda)\Phi_c = \lambda\Phi_c, \quad (1)$$

where $\lambda\Phi_c = (\lambda\phi_{1c}, \lambda\phi_{2c})^\top$. Since $L_{\Phi_c}(\lambda)$ are differential operators in the variables x_j while λ depends only on z , (1) implies the commutation condition

$$L_{\Phi_c}(\lambda\mu) = L_{\Phi_c}(\lambda)L_{\Phi_c}(\mu) = L_{\Phi_c}(\mu)L_{\Phi_c}(\lambda),$$

where $\mu(z) \in \mathcal{A}_\Theta$. We thus arrive at the following

Corollary [1]. *There is an embedding of the rings*

$$L_{\Phi_c} : \mathcal{A}_\Theta \rightarrow \text{Mat}(2, \mathcal{D}),$$

where $\text{Mat}(2, \mathcal{D})$ is the ring of (2×2) -matrix differential operators. The range of the embedding is a commutative ring of differential operators.

Now, we suppose that Θ is a nonsingular Riemann surface.

PROOF OF THEOREM 1. We start with proving the second part of the theorem. Introduce the following functions in M_c :

$$\begin{aligned} \psi &= \frac{\theta(z + c + x)}{\theta(z)} \exp(-x_1 \partial_{z_1} \log \theta(z) - x_2 \partial_{z_2} \log \theta(z)), \\ \psi_{c'} &= \frac{\theta(z + c + c' + x) \theta(z - c')}{\theta^2(z)} \exp(-x_1 \partial_{z_1} \log \theta(z) - x_2 \partial_{z_2} \log \theta(z)). \end{aligned}$$

They determine a basis for the $\mathcal{D}$ -module M_c [3]. In [3] we have found the Nakayashiki operators $L_{c,c'}(\mathcal{A}_\Theta)$ in this basis. The coefficients of the operators $L_{c,c'}(\lambda)$ are doubly periodic; moreover, the coefficients of all components of each operator have singularity at $x = -c$ [3]. Let X be the Jacobi variety of a Riemann surface with real branching points. In this case we have $\bar{\theta}(z) = \theta(\bar{z})$ [5]. Suppose also that $c, c' \in \mathbb{R}^2$ and $\lambda(z)$ is a real function for $z \in \mathbb{R}^2$ (for example, $\lambda = \partial_{z_j} \partial_{z_i} \log \theta(z)$). The fact that the operators $L_{c,c'}(\lambda)$ are real is immediate from the fact that the eigenfunction $(\psi, \psi_{c'})^\top$ of the operator $L_{c,c'}(\lambda)$ is real for real values of z .

We turn to proving the first part of the theorem. First of all, observe that M_0 is not a free $\mathcal{D}$ -module of rank 2. This follows, for example, from the fact that $\partial_{x_j}\psi(z, 0) = 0$ which implies that a function of the form $f(z, x) \exp(-x_1 \partial_{z_1} \log \theta(z) - x_2 \partial_{z_2} \log \theta(z)) \in M_0$, where $f(z, x)$ has a pole of the second order on the theta-divisor, cannot in general be represented as $d_1 \psi + d_2 \psi_{c'}$ for $x = 0$, where $d_1 \in \mathcal{D}$ is a first-order operator and d_2 is the operator of multiplication by a function; however, this is possible for $x \neq 0$.

Let X be the Jacobi variety of the Riemann surface Γ ; $L(\lambda)$, a Nakayashiki operator with doubly periodic coefficients with periods $\tau_1, \tau_2 \in \mathbb{R}$; $\Psi = (\psi_1, \psi_2)^\top$, a Bloch eigenfunction whose components are basis elements of the $\mathcal{D}$ -module M_c ; and

$$\begin{aligned} L(\lambda)\Psi &= \lambda\Psi, \quad \Psi(z, x_1 + \tau_1, x_2) = -\partial_{z_1} \log \theta(z) \tau_1 \mu_1 \Psi(z, x_1, x_2), \\ \Psi(z, x_1, x_2 + \tau_2) &= -\partial_{z_2} \log \theta(z) \tau_2 \mu_2 \Psi(z, x_1, x_2), \end{aligned}$$

where $\mu_1, \mu_2 \in \mathbb{C}$. Suppose that $L(\lambda)$ has real coefficients for $x \in \mathbb{R}^2$. Then X admits an antiholomorphic involution τ such that

$$\overline{\Psi}(z, x) = \Psi(\tau(z), x), \quad \bar{\lambda}(z) = \lambda(\tau(z)).$$

Moreover, τ leaves the theta-divisor invariant, $\tau(\Theta) = \Theta$, since the multiplicative functions $\partial_{z_1} \log \theta(z)$ and $\partial_{z_2} \log \theta(z)$ have a pole on the theta-divisor. Consequently, τ is induced by an antiholomorphic involution of Γ and coincides with complex conjugation. The theta-divisor is invariant under τ only if Γ has real branching points [5].

Let us show that $c \in \mathbb{R}^2$. Denote the transition operator from the basis $\psi, \psi_{c'}$ to the basis ψ_1, ψ_2 by A . The equalities $\Psi = A(\psi, \psi_{c'})^\top$, $\bar{\theta}(z) = \theta(\bar{z})$, and $\Psi(z + \Omega m, x) = \exp(-2\pi i \langle m, c \rangle) \Psi(z, x)$ then imply that $\overline{\Psi}(z + \Omega m, x) = \exp(-2\pi i \langle m, \bar{c} \rangle) \overline{\Psi}(\bar{z}, x)$. Hence, from $\Psi(z, x) = \overline{\Psi}(\bar{z}, x)$ we obtain $c \in \mathbb{R}^2$.

The cause for nonsmoothness of the operators is as follows. Replace $x + c$ with x and divide ψ_1 and ψ_2 by $\exp(-c_1 \partial_{z_1} \log \theta(z) - c_2 \partial_{z_2} \log \theta(z))$ to obtain the functions $\tilde{\psi}_1, \tilde{\psi}_2 \in M_0$. Since the $\mathcal{D}$ -module M_0 is not free of rank 2, the equality $\tilde{L}(\lambda)\tilde{\Psi} = \lambda\tilde{\Psi}$, $\tilde{\Psi} = (\tilde{\psi}_1, \tilde{\psi}_2)^\top$, does not hold in a neighborhood of $x = 0$. Consequently, the coefficients of $\tilde{L}(\lambda)$ have singularity at $x = 0$, and so the coefficients of $L(\lambda)$ have singularity at $x = -c$. Namely, if the coefficients of A and A^{-1} are smooth at $x = -c$ then the equality $A^{-1}L(\lambda)A = L_{c,c'}(\lambda)$ [3] implies that the operator $L(\lambda)$ is nonsmooth at $x = -c$. If A or A^{-1} is nonsmooth at $x = -c$ then nonsmoothness of $L(\lambda)$ results from the equality

$L(\lambda) = AL_{c,c'}(\lambda)A^{-1}$ and the fact that all components of the operator $L_{c,c'}(\lambda)$ are nonsmooth for $x = -c$. Theorem 1 is proven.

Denote by Γ the smooth completion of the Riemann surface that is defined in the (y, w) -plane by the equation $w^2 = P(y) = (y - y_1) \dots (y - y_5)$ with real $y_1 < \dots < y_5$. Denote by X the Jacobi variety of the hyperelliptic surface Γ . The holomorphic involution $\sigma : (y, w) \rightarrow (y, -w)$ acts on Γ with the fixed points $Q_i = (y_i, 0), i = 1, \dots, 5, \infty$, and we also have the antiholomorphic involution $\tau : (y, w) \rightarrow (\bar{y}, \bar{w})$. The involution τ has three fixed cycles:

$$C_1 : \{y_1 \leq y \leq y_2, w = \pm \sqrt{P(y)}\}, \quad C_2 : \{y_3 \leq y \leq y_4, w = \pm \sqrt{P(y)}\},$$

$$C_3 : \{y_5 \leq y \leq \infty, w = \pm \sqrt{P(y)}\}.$$

Take a canonical basis a_1, a_2, b_1, b_2 of cycles with the intersection indices $a_i \circ a_j = b_i \circ b_j = 0$ and $a_i \circ b_j = \delta_{ij}$ as shown in Fig. 1 (the dots indicate parts of cycles on the “lower leaf” of the Riemann surface). The union of the cycles C_1, C_2 , and C_3 divides Γ into two disjoint pieces.

Observe that $a_1 = C_1$ and $a_2 = C_2$. The antiholomorphic and holomorphic involutions act on these cycles as follows:

$$\tau a_1 = a_1, \quad \tau a_2 = a_2, \quad \tau b_1 = -b_1, \quad \tau b_2 = -b_2,$$

$$\sigma a_1 = -a_1, \quad \sigma a_2 = -a_2, \quad \sigma b_1 = -b_1, \quad \sigma b_2 = -b_2,$$

where equality is understood to be in the homology group. On the corresponding canonical basis of abelian differentials ω_1 and ω_2 such that

$$\int_{a_j} \omega_i = \delta_{ij}, \quad \int_{b_j} \omega_i = \int_{b_i} \omega_j = \Omega_{ij}, \quad i, j = 1, 2,$$

the involutions τ and σ act as follows:

$$\tau^* \omega_i = \bar{\omega}_i, \quad \sigma^* \omega_i = -\omega_i. \tag{2}$$

Introduce four real tori in X :

$$T_1 : z \equiv i(t_1, t_2), \quad T_2 : z \equiv i(t_1, t_2) + \left(\frac{1}{2}, 0\right),$$

$$T_3 : z \equiv i(t_1, t_2) + \left(0, \frac{1}{2}\right), \quad T_4 : z \equiv i(t_1, t_2) + \left(\frac{1}{2}, \frac{1}{2}\right),$$

where $(t_1, t_2) \in \mathbb{R}^2$, the symbol $\equiv$ stands for equality modulo an element of the lattice $\mathbb{Z}^2 + \Omega\mathbb{Z}^2$, and the matrix Ω is composed of the entries Ω_{ij} . The theta-function is real on these tori.

Lemma 2. *The following equalities are valid:*

$$\begin{aligned} \int_{\infty}^{Q_1} \omega &\equiv \left(\frac{\Omega_{11}}{2}, \frac{\Omega_{12}}{2} \right), \quad \int_{\infty}^{Q_2} \omega \equiv \left(\frac{1}{2} + \frac{\Omega_{11}}{2}, \frac{\Omega_{12}}{2} \right), \\ \int_{\infty}^{Q_3} \omega &\equiv \left(\frac{1}{2} - \frac{\Omega_{21}}{2}, -\frac{\Omega_{22}}{2} \right), \quad \int_{\infty}^{Q_4} \omega \equiv \left(\frac{1}{2} - \frac{\Omega_{21}}{2}, \frac{1}{2} - \frac{\Omega_{22}}{2} \right), \quad \int_{\infty}^{Q_5} \omega \equiv \left(\frac{1}{2}, \frac{1}{2} \right). \end{aligned}$$

PROOF Join ∞ and Q_1 by an oriented path l such that $l \cup -\sigma l = b_1$ ($-\sigma l$ stands for σl with opposite orientation). Then

$$\int_l \omega - \int_{\sigma l} \omega = \int_{b_1} \omega, \quad \int_l \omega = \int_{\sigma l} \sigma^* \omega = - \int_{\sigma l} \omega;$$

consequently,

$$\int_{\infty}^{Q_1} \omega = \frac{1}{2} \int_{b_1} \omega \equiv \left(\frac{\Omega_{11}}{2}, \frac{\Omega_{12}}{2} \right).$$

Similarly, we demonstrate that

$$\begin{aligned} \int_{Q_1}^{Q_2} \omega &\equiv \left(\frac{1}{2}, 0 \right), \quad \int_{Q_2}^{Q_3} \omega = \frac{1}{2} \int_{b_1 - b_2} \omega \equiv \left(\frac{\Omega_{11}}{2} - \frac{\Omega_{21}}{2}, \frac{\Omega_{12}}{2} - \frac{\Omega_{22}}{2} \right), \\ \int_{Q_3}^{Q_4} \omega &= \frac{1}{2} \int_{a_2} \omega \equiv \left(0, \frac{1}{2} \right), \quad \int_{Q_4}^{Q_5} \omega = \frac{1}{2} \int_{b_2} \omega \equiv \left(\frac{\Omega_{21}}{2}, \frac{\Omega_{22}}{2} \right). \end{aligned}$$

The lemma is proven.




Fig. 1.

As shown in [7], with this choice of a canonical basis of cycles and the choice of ∞ as the initial point of the Abel mapping, the vector of Riemann constants equals

$$K \equiv \left(\frac{\Omega_{11}}{2} + \frac{\Omega_{12}}{2}, \frac{\Omega_{21}}{2} + \frac{\Omega_{22}}{2} \right) + \left(1, \frac{1}{2} \right).$$

Lemma 3. *The theta-function $\theta(z)$ has no zeros on T_1 .*

PROOF. Suppose that z belongs to T_1 and to the theta-divisor. Then $\bar{z} \equiv -z$ and, by the Riemann theorem about the zeros of a theta-function (see [8]), $z \equiv A(P) + K$, where $A(P)$ is the Abel mapping with the initial point ∞ . From (2) and the fact that Ω is a purely imaginary matrix we obtain the equalities

$$\begin{aligned} \bar{z} &\equiv A(\tau(P)) - \left(\frac{\Omega_{11}}{2} + \frac{\Omega_{12}}{2}, \frac{\Omega_{21}}{2} + \frac{\Omega_{22}}{2} \right) + \left(1, \frac{1}{2} \right) \equiv -z \equiv \\ &-A(P) - \left(\frac{\Omega_{11}}{2} + \frac{\Omega_{12}}{2}, \frac{\Omega_{21}}{2} + \frac{\Omega_{22}}{2} \right) - \left(1, \frac{1}{2} \right); \end{aligned}$$

consequently, $A(\tau(P)) + A(P) \equiv 0$. Together with (2), this yields $\tau(P) = \sigma(P)$. Hence, either P is a branching point or the y -coordinate of P is real and the w -coordinate of P is purely imaginary. These points constitute the three cycles

$$\begin{aligned} B_1 &: \{ \infty \leq y \leq y_1, w = \pm \sqrt{P(y)} \}, \quad B_2 : \{ y_2 \leq y \leq y_3, w = \pm \sqrt{P(y)} \}, \\ B_3 &: \{ y_4 \leq y \leq y_5, w = \pm \sqrt{P(y)} \}. \end{aligned}$$

For the points of these cycles we have

$$\overline{A(P)} \equiv A(\tau(P)) \equiv A(\sigma(P)) \equiv -A(P).$$

Hence,

$$A(P) \equiv i(t_1, t_2) + \frac{(m, n)}{2}, \quad (t_1, t_2) \in \mathbb{R}^2, \quad m, n \in \mathbb{Z}.$$

Consequently, the real part of $A(P)$ does not change upon the circuit around the cycles B_j , $j = 1, 2, 3$. Since $Q_1 \in B_1$, $Q_2 \in B_2$, and $Q_5 \in B_3$, Lemma 2 implies the inclusions $A(B_1) \subset T_1$, $A(B_2) \subset T_2$, and $A(B_3) \subset T_4$. Therefore, $z \equiv A(P) + K$ cannot belong to T_1 . The lemma is proven.

Put

$$c' \equiv \left(\frac{\Omega_{11}}{2} - \frac{\Omega_{21}}{2}, \frac{\Omega_{12}}{2} - \frac{\Omega_{22}}{2} \right).$$

Lemma 4. *The theta-divisor and the Riemann surface defined in X by the equation $\theta(z - c') = 0$ intersect at the two points*

$$p_1 \equiv \left(\frac{\Omega_{12}}{2} + \frac{1}{2}, \frac{\Omega_{22}}{2} + \frac{1}{2} \right), \quad p_2 \equiv \left(\frac{\Omega_{11}}{2} + \frac{1}{2}, \frac{\Omega_{21}}{2} + \frac{1}{2} \right).$$

PROOF. A point z belongs to the theta-divisor whenever it has the form $z \equiv \int_{\infty}^P \omega + K$, where $P \in \Gamma$. Consequently, the intersection points look like

$$p_1 \equiv \int_{\infty}^{P_1} \omega + K, \quad p_2 \equiv \int_{\infty}^{P_2} \omega + K,$$

where P_1 and P_2 are zeros of the function $\theta(\int_{\infty}^P \omega + K - c')$ on Γ . The function $\theta(\int_{\infty}^P \omega + K - c')$ is not identically zero on Γ (since $K - c' \equiv (1, \frac{1}{2}) \not\equiv K$). Then by the Riemann theorem

$$\int_{\infty}^{P_1} \omega + \int_{\infty}^{P_2} \omega \equiv c'$$

and the points P_1 and P_2 are determined uniquely by c' (see, for example, [7]). By Lemma 2,

$$\int_{\infty}^{Q_2} \omega + \int_{\infty}^{Q_3} \omega \equiv c';$$

consequently,

$$p_1 \equiv \int_{\infty}^{Q_2} \omega + K, \quad p_2 \equiv \int_{\infty}^{Q_3} \omega + K.$$

The lemma is proven.

From now on, we assume that $c \in T_1$. Henceforth we use the following notations:

$$\theta_j(z) = \partial_{z_j} \theta(z), \quad \theta_{kj}(z) = \partial_{z_k} \partial_{z_j} \theta(z), \quad k, j = 1, 2.$$

PROOF OF THEOREM 2. To find smooth real operators, we must make some modifications in the formulas of [3] where, proving Propositions 2 and 3, we used the fact that $\theta(\Delta) = 0$, where Δ stands for the vector of Riemann constants. The other properties of Δ are not used in the proof of these propositions. All formulas for the Nakayashiki operators of Propositions 2 and 3 of [3] remain valid if we replace Δ with $p_3 = \int_{\infty}^{P_3} \omega + K$, where P_3 is an arbitrary point of B_2 other than Q_2 or Q_3 and $\theta(p_3) = 0$, $p_3 \in T_4$. Calculating $[L_{c,c'}(\partial_{z_k} \partial_{z_j} \log \theta(z))]_{12}$ in Proposition 2 of [3], we put $z = 0$. The formula remains valid if we put $z = p_4$, where p_4 is an arbitrary point of T_4 such that $\theta(p_4) \neq 0$. Then

$$\begin{aligned} [L_{c,c'}(\partial_{z_k} \partial_{z_j} \log \theta(z))]_{12} &= \frac{\theta^2(p_4)}{\theta(p_4 + c + c' + x)\theta(c')} \\ &\times (\partial_{z_k} \partial_{z_j} - f_{c,c'}^{kj} \partial_{z_1} - g_{c,c'}^{kj} \partial_{z_2} - h_{c,c'}^{kj} + 2\partial_{z_k} \partial_{z_j} \log \theta(z)) \left(\frac{\theta(z + c + x)}{\theta(z)} \right) \Big|_{z=p_4}. \end{aligned}$$

Similarly, in the formula for $g_{c,c'}^j$ of Proposition 3 we now put $z = p_4$ instead of $z = 0$. It is these formulas that we will use below.

The constants $\theta_k(p_j)$ are imaginary and the functions $\partial_{x_k} \log \theta(p_j + c + x)$ are purely imaginary for $x \in i\mathbb{R}^2 + (\frac{1}{2}, \frac{1}{2})$; consequently, the functions $f_{c,c'}^{kj}$ and $g_{c,c'}^{kj}$ in Proposition 2 of [3] are purely imaginary and $h_{c,c'}^{kj}$ are real. Hence, the operators $[L_{c,c'}(\partial_{z_j} \partial_{z_k} \log \theta(z))]_{11}$ and $[L_{c,c'}(\partial_{z_j} \partial_{z_k} \log \theta(z))]_{12}$ are real. The numbers α_{kj} and α in Proposition 2 of [3] are real, since the functions $\partial_{z_j} \partial_{z_k} \log \theta(z)$ and $\frac{\theta(z-c')\theta(z+c')}{\theta^2(z)}$ are real on T_1 and the functions $\partial_{z_j} \partial_{z_k} \log \theta(z)$ are linearly independent (this follows, for example, from Nakayashiki's theorem, since the 11-coefficients of the operators $L_{c,c'}(\partial_{z_j} \partial_{z_k} \log \theta(z))$ are equal to $\partial_{x_j} \partial_{x_k}$). Consequently, the operators

$$[L_{c,c'}(\partial_{z_j} \partial_{z_k} \log \theta(z))]_{21}, \quad [L_{c,c'}(\partial_{z_j} \partial_{z_k} \log \theta(z))]_{22}$$

are real.

Since the coefficients of the Nakayashiki operators are expressed in terms of the theta-function and its derivatives while the theta-function is real-valued on T_1 and T_4 , the fact that the coefficients of the operators are real

for $x \in i\mathbb{R}^2$ implies that so are the coefficients for $x \in i\mathbb{R}^2 + (\frac{1}{2}, \frac{1}{2})$. From Proposition 3 of [3] we infer that the coefficients of the operators Z_1 and Z_2 [2] are purely imaginary for $x \in i\mathbb{R}^2$. Then Proposition 1 implies that the operators $L_{c,c'}(i\partial_{z_s}\partial_{z_j}\partial_{z_k}\log\theta(z))$ are real for $x \in i\mathbb{R}^2$.

By Propositions 1, 2, and 3 of [3] with the above-made modifications, to prove smoothness we have to demonstrate that the functions $\theta(p_1 + c + x)$, $\theta(p_2 + c + x)$, $\theta(p_3 + c + c' + x)$, and $\theta(p_4 + c' + c + x)$ do not vanish for $x \in i\mathbb{R}^2 + (\frac{1}{2}, \frac{1}{2})$. This follows from Lemma 3.

Introduce the magnetic translation operators $\tilde{T}_1$ and $\tilde{T}_2$:

$$\tilde{T}_1\varphi(x) = \varphi(x + \Omega_1)\exp(2\pi i x_1), \quad \tilde{T}_2\varphi(x) = \varphi(x + \tilde{\Omega}_2)\exp(2\pi i x_2),$$

where Ω_j is the j th row of Ω . The functions ψ and $\psi_{c'}$ are magneto-Bloch functions:

$$\tilde{T}_1\psi = \mu_1\psi, \quad \mu_1 = \exp(-\pi i\Omega_{11} - 2\pi i(z_1 + c_1) - \Omega_{11}\partial_{z_1}\log\theta(z) - \Omega_{12}\partial_{z_2}\log\theta(z)),$$

$$\tilde{T}_2\psi = \mu_2\psi, \quad \mu_2 = \exp(-\pi i\Omega_{22} - 2\pi i(z_2 + c_2) - \Omega_{12}\partial_{z_1}\log\theta(z) - \Omega_{22}\partial_{z_2}\log\theta(z)),$$

$$\tilde{T}_1\psi_{c'} = \mu_1\exp(-2\pi i c'_1)\psi_{c'}, \quad \tilde{T}_2\psi_{c'} = \mu_2\exp(-2\pi i c'_2)\psi_{c'}.$$

Instead of $\psi_{c'}$ we use the function

$$\tilde{\psi}_{c'} = \psi_{c'}\exp\left(2\pi i\left\langle x - \left(\frac{1}{2}, \frac{1}{2}\right), \Omega^{-1}c' \right\rangle\right).$$

Then by symmetry of Ω

$$\tilde{T}_1\tilde{\psi}_{c'} = \mu_1\tilde{\psi}_{c'}, \quad \tilde{T}_2\tilde{\psi}_{c'} = \mu_2\tilde{\psi}_{c'}.$$

The Nakayashiki operators in the basis ψ , $\tilde{\psi}_{c'}$ look like $dL_{c,c'}(\lambda)d^{-1}$, where d is the diagonal matrix with diagonal $(1, \exp(2\pi i\langle x - (\frac{1}{2}, \frac{1}{2}), \Omega^{-1}c' \rangle))$ [3], and are smooth and real under the same conditions as the operators $L_{c,c'}(\lambda)$. Denote by H the smooth operator $dL_{c,c'}(\partial_{z_1}^2\log\theta(z) + \partial_{z_2}^2\log\theta(z))d^{-1}$ which is real for $x \in i\mathbb{R}^2 + (\frac{1}{2}, \frac{1}{2})$. Its 11-component is

$$H_{11} = (i\partial_{x_1} - A_1)^2 + (i\partial_{x_2} - A_2)^2 + u(x),$$

where [3]

$$A_1 = \frac{i}{2}(f_{c,c'}^{11} + f_{c,c'}^{22}), \quad A_2 = \frac{i}{2}(g_{c,c'}^{11} + g_{c,c'}^{22}).$$

This is the Schrödinger operator in the periodic magnetic field $rmrot(A_1, A_2, 0)$. For the components of the vector-potential (A_1, A_2) we have the equality

$$A_k(x + \Omega_j) - A_k(x) = 2\pi\delta_{kj}.$$

The magnetic translation operators commute with the covariant differentiation operators:

$$\tilde{T}_j(i\partial_{x_k} - A_k) = (i\partial_{x_k} - A_k)\tilde{T}_j.$$

The magneto-Bloch function ψ for $z \in \Gamma_{\mathcal{C}'}$ satisfies the Schrödinger equation with the Hamiltonian H_{11} and the energy $\partial_{z_1}^2 \log \theta(z) + \partial_{z_2}^2 \log \theta(z)$. Hence, the potential is doubly periodic; i.e., $u(x + \Omega_1) = u(x + \Omega_2) = u(x)$, and the Schrödinger operator commutes with the magnetic translation operators: $\tilde{T}_j H_{11} = H_{11} \tilde{T}_j$. The 12-component of H is the operator of multiplication by a doubly periodic function. The operator H_{21} looks like $F(x)\tilde{H}_{21}$, where F is a doubly periodic function and $\tilde{H}_{21}$ is a second-order operator with constant leading coefficients. The 22-component is as follows:

$$H_{22} = (i\partial_{x_1} - \tilde{A}_1)^2 + (i\partial_{x_2} - \tilde{A}_2)^2 + \tilde{u}(x).$$

The operator H_{22} possesses the same properties as H_{11} . In particular, H_{22} commutes with the magnetic translation operators.

The magneto-Bloch functions of H at an energy level λ are parametrized by the Riemann surface defined in X by the equation

$$\partial_{z_1}^2 \log \theta(z) + \partial_{z_2}^2 \log \theta(z) = \lambda.$$

To complete the proof, we have to change coordinates and recall that ∂_{x_k} are the operators of complex differentiation; i.e., $\partial_{x_k} = \frac{1}{2}(\frac{\partial}{\partial \bar{x}_k} - i\frac{\partial}{\partial y_k})$.

Theorem 2 is proven.

3 The Nakayashiki Operators

In the beginning of this section we introduce two Riemann surfaces Γ_1 and Γ_2 of genus 2 embedded in the two-dimensional abelian variety X . In Lemma 5, we use the Fay formula (3) to prove that Γ_1 and Γ_2 are tangent to the theta-divisor. In (5) and (6), we indicate a basis ψ_1, ψ_2 for the Baker–Akhiezer module. It follows from Lemma 5 that ψ_1 and ψ_2 , bounded on Γ_1 and Γ_2 respectively, are one-point Baker–Akhiezer functions [9]. In Lemma 7

we find some coefficients of the 11- and 12-components of the second-order operators. In Lemma 8 we indicate a connection between the 11- and 12-components and between the 21- and 22-components of the operators. In Lemma 9 we find some coefficients of L_1 . Lemma 1 ensues from Lemmas 7–9. In Lemma 10 we prove that the coefficients of the 11-components of the Nakayashiki operators are rationally expressible in terms of the function V and its derivatives. Theorem 3 ensues from Lemmas 11 and 12. In the latter we prove that the coefficients of the operators of the second and third orders are rationally expressible in terms of V , W , and their derivatives.

The abelian variety X is the Jacobi variety of some Riemann surface Γ of genus 2. There is a canonical basis a_1, a_2, b_1, b_2 of cycles on Γ with the intersection indices $a_i \circ a_j = b_i \circ b_j = 0$ and $a_i \circ b_j = \delta_{ij}$, and there is a basis of abelian differentials ω_1 and ω_2 such that $\int_{a_j} \omega_i = \delta_{ij}$, $i, j = 1, 2$, and the components of Ω are equal to $\Omega_{ij} = \int_{b_j} \omega_i$. The following identity due to J. D. Fay [10] is valid for points $\tilde{R}, \tilde{Q} \in \Gamma$:

$$\sum_{i,j=1}^2 F_i G_j \partial_{z_i} \partial_{z_j} \log \theta(z) = \tilde{c}_3 + \tilde{c}_2 \frac{\theta\left(z + \int_{\tilde{Q}}^{\tilde{R}} \omega\right) \theta\left(z + \int_{\tilde{R}}^{\tilde{Q}} \omega\right)}{\theta^2(z)},$$

where $F_i = \frac{\omega_i(\tilde{R})}{dr}$, $G_i = \frac{\omega_i(\tilde{Q})}{dq}$, r and q are local parameters in neighborhoods of $\tilde{R}$ and $\tilde{Q}$,

$$\int_{\tilde{Q}}^{\tilde{R}} \omega = \left(\int_{\tilde{Q}}^{\tilde{R}} \omega_1, \int_{\tilde{Q}}^{\tilde{R}} \omega_2 \right),$$

and $\tilde{c}_2$ and $\tilde{c}_3$ are some constants. Denote by $R, Q \in \Gamma$ the zeros of ω_2 . Since $\int_Q^R \omega = -2K$, where K is the vector of Riemann constants with respect to Q (see, for example, [11]), we have

$$\partial_{z_1}^2 \log \theta(z) = c_3 + c_2 \frac{\theta(z - 2K) \theta(z + 2K)}{\theta^2(z)}, \quad (3)$$

where c_2 and c_3 are some constants (we write them down explicitly in (15)). Denote by Γ_1 and Γ_2 the Riemann surfaces that are defined in X by the equations $\theta(z + 2K) = 0$ and $\theta(z - 2K) = 0$.

Lemma 5. *The Riemann surfaces Θ and Γ_1 (Θ and Γ_2) intersect at the point $-K$ (K) with multiplicity 2 (are tangent), and $\theta_{11}(-K) = \theta_{11}(K) \neq 0$.*

PROOF. The index of intersection of Θ and Γ_1 equals 2 (see [8]). Suppose that Θ and Γ_1 intersect at two different points. Let $A : \Gamma \rightarrow X$ be the Abel mapping given by the formula

$$A(P) = \left(\int_Q^P \omega_1, \int_Q^P \omega_2 \right), \quad P \in \Gamma.$$

By the Riemann theorem about the zeros of a theta-function, the equality $\theta(z) = 0$ amounts to the fact that $z = A(P) + K$. Consequently, $z = A(R) + K = -K$ is the intersection point of Θ and Γ_1 . Take a local parameter s at the point R . It follows from (3) that $\theta_1(K) = 0$. The following equality holds:

$$\frac{d}{ds} \theta \left(\int_{s(Q)}^{s(P)} s^* \omega + 3K \right) = \theta_1 \left(\int_{s(Q)}^{s(P)} s^* \omega + 3K \right) \frac{\omega_1}{ds} + \theta_2 \left(\int_{s(Q)}^{s(P)} s^* \omega + 3K \right) \frac{\omega_2}{ds}.$$

Since $\omega_2(R) = 0$ and $\theta_1(K) = 0$, we have

$$\frac{d}{ds} \theta \left(\int_{s(Q)}^{s(P)} s^* \omega + 3K \right) = 0$$

for $P = R$; consequently, the function $\theta(A(P) + K + 2K)$ has zero of multiplicity 2 at R or, equivalently, the function $\theta(z + 2K)$ has zero of multiplicity 2 on Θ at $-K$. Hence, the point $-K$ is a tangency point of Θ and Γ_1 . Similarly, we can prove that Θ and Γ_2 are tangent at K .

Prove that $\theta_{11}(-K) \neq 0$. From (3) we obtain the identity

$$\begin{aligned} & \theta_{112}(z)\theta(z) - \theta_{11}(z)\theta_2(z) - 2\theta_{12}(z)\theta_1(z) - 2c_3\theta_2(z)\theta(z) \\ &= c_2\theta_2(z + 2K)\theta(z - 2K) + c_2\theta(z + 2K)\theta_2(z - 2K) \end{aligned}$$

which implies that $\theta_{11}(-K) \neq 0$; otherwise we would have $\theta_2(K) = 0$, but Θ is a smooth Riemann surface. The lemma is proven.

From (3) we conclude that the following equality is valid on Γ_1 and Γ_2 :

$$\theta_{11}(z)\theta(z) - \theta_1^2(z) - c_3\theta^2(z) = 0.$$

Lemma 5 implies that the function $\theta_1(z)$ on Γ_1 and Γ_2 has zeros of the first order at $-K$ and K ; consequently, we have the expansions

$$\theta_2(z) = \theta_2(-K) + b_1\theta_1(z) + o(\theta_1(z)), \quad \theta_{11}(z) = \theta_{11}(-K) + d_1\theta_1(z) + o(\theta_1(z)) \quad (4)$$

on Γ_1 (in a neighborhood of the point $-K \in \Gamma_1$) and the expansions

$$\theta_2(z) = \theta_2(K) + b_2\theta_1(z) + o(\theta_1(z)), \quad \theta_{11}(z) = \theta_{11}(K) + d_2\theta_1(z) + o(\theta_1(z))$$

on Γ_2 (in a neighborhood of the point $K \in \Gamma_2$), where $b_i, d_i \in \mathbb{C}$, $i = 1, 2$.

Lemma 6. *The equalities $b_1 = b_2$ and $d_1 = -d_2$ hold.*

PROOF. Denote by A_1 the mapping $\Gamma \rightarrow X$ which is defined by the formula $A_1(P) = \int_Q^P \omega - K$, $P \in \Gamma$, and suppose that $A_2(P) = -A_1(P)$. The range of A_1 is the Riemann surface Γ_1 and the range of A_2 is Γ_2 . Since $\theta(z)$ is an even function, $\theta_1(z)$ and $\theta_2(z)$ are odd; consequently, from (4) we obtain

$$\theta_2(A_2(P)) = \theta_2(K) + b_1\theta_1(A_2(P)) + \dots$$

Hence, $b_1 = b_2$. Similarly, we can prove that $d_1 = -d_2$. The lemma is proven.

Put $b = b_1 = b_2$ and $d = d_1 = -d_2$.

Introduce the following functions in M_c :

$$\begin{aligned} \psi_1 &= \frac{\theta(z + c + x)}{\theta(z)\theta(c - K + x)} \\ &\times \exp \left(-x_1 \left(\partial_{z_1} \log \theta(z) - \frac{b\theta_{11}(K)}{2\theta_2(K)} - \frac{d}{2} \right) - x_2 \partial_{z_2} \log \theta(z) \right), \end{aligned} \quad (5)$$

$$\begin{aligned} \psi_2 &= \frac{\theta(z + c - 2K + x)\theta(z + 2K)}{\theta^2(z)\theta(c - K + x)} \\ &\times \exp \left(-x_1 \left(\partial_{z_1} \log \theta(z) + \frac{b\theta_{11}(K)}{2\theta_2(K)} + \frac{d}{2} \right) - x_2 \partial_{z_2} \log \theta(z) \right). \end{aligned} \quad (6)$$

The functions ψ_1 and ψ_2 constitute a basis for the $\mathcal{D}$ -module M_c [3].

Denote by $L_{c,K}^{ij}$ and $L_{c,K}^{ijk}$ the Nakayashiki operators $L_{c,K}(\partial_{z_i}\partial_{z_j} \log \theta(z))$ and $L_{c,K}(\partial_{z_i}\partial_{z_j}\partial_{z_k} \log \theta(z))$ in the basis ψ_1, ψ_2 (the meaning of the subscript K will be seen later). The operator $[L_{c,K}^{ij}]_{11}$ looks like $-\partial_{x_i}\partial_{x_j} + f_{c,K}^{ij}\partial_{x_1} +$

$g_{c,K}^{ij}\partial_{x_2} + h_{c,K}^{ij}$ and the operator $[L_{c,K}(\partial_{z_i}\partial_{z_j}\log\theta(z))]_{12}$ is the operator of multiplication by some function $H_{c,K}^{ij}(x)$, $i, j = 1, 2$ [3].

Lemma 7. *The following equalities hold:*

$$g_{c,K}^{11} = \frac{\theta_{11}(K)}{\theta_2(K)}, \quad g_{c,K}^{12} = \frac{b\theta_{11}(K)}{2\theta_2(K)} + \frac{\theta_{12}(K)}{\theta_2(K)} + \frac{d}{2}, \quad g_{c,K}^{22} = \frac{\theta_{22}(K)}{\theta_2(K)}, \quad f_{c,K}^{11} = 0,$$

$$\begin{aligned} h_{c,K}^{11} &= \partial_{x_1}^2 \log \frac{\theta(c-3K+x)}{\theta(c-K+x)} \\ &+ \left(\partial_{x_1} \log \frac{\theta(c-3K+x)}{\theta(c-K+x)} + \partial_{z_1} \log \theta(3K) + \frac{b\theta_{11}(K)}{2\theta_2(K)} + \frac{d}{2} \right)^2 \\ &- \frac{\theta_{11}(K)}{\theta_2(K)} \left(\partial_{x_2} \log \frac{\theta(c-3K+x)}{\theta(c-K+x)} + \partial_{z_2} \log \theta(3K) \right) + c_3, \\ H_{c,K}^{11} &= \frac{2\theta_{11}(K)\theta(K+c+x)}{\theta(3K)\theta(c-K+x)} \exp(x_1(\frac{b\theta_{11}(K)}{\theta_2(K)} + d)), \\ H_{c,K}^{12} &= \left(\frac{\theta_{12}(K)}{\theta_{11}(K)} + \frac{b}{2} + \frac{d\theta_2(K)}{2\theta_{11}(K)} \right) H_{c,K}^{11} - \frac{\theta_2(K)}{2\theta_{11}(K)} \partial_{x_1} H_{c,K}^{11}, \\ H_{c,K}^{22} &= \left(\frac{\theta_{22}(K)}{\theta_{11}(K)} + b + \frac{d\theta_2(K)}{\theta_{11}(K)} \right) H_{c,K}^{11} - \frac{\theta_2(K)}{\theta_{11}(K)} \partial_{x_2} H_{c,K}^{11}. \end{aligned}$$

PROOF. Divide the equality

$$-\partial_{x_i}\partial_{x_j}\psi_1 + f_{c,K}^{ij}\partial_{x_1}\psi_1 + g_{c,K}^{ij}\partial_{x_2}\psi_1 + h_{c,K}^{ij}\psi_1 + H_{c,K}^{ij}\psi_2 = \partial_{z_i}\partial_{z_j}\log\theta(z)\psi_1$$

by $\exp(-x_1\partial_{z_1}\log\theta(z) - x_2\partial_{z_2}\log\theta(z))$ and multiply by $\theta^2(z)$. Putting first $z = -K$ and then $z = K$, we obtain $g_{c,K}^{ij}$ and $H_{c,K}^{ij}$. Now, take $z \in \Gamma_1$. Using (4), we find that $f_{c,K}^{11} = 0$. The following equality is valid for $z \in \Gamma_1$:

$$\partial_{x_1}^2 \psi_1 - \frac{\theta_{11}(K)}{\theta_2(K)} \partial_{x_2} \psi_1 - h_{c,K}^{11} \psi_1 + c_3 \psi_1 = 0; \quad (7)$$

consequently,

$$h_{c,K}^{11} = \partial_{x_1}^2 \log \psi_1 + (\partial_{x_1} \log \psi_1)^2 - \frac{\theta_{11}(K)}{\theta_2(K)} \partial_{x_2} \log \psi_1 + c_3.$$

Putting $z = -3K \in \Gamma_1$, we obtain $h_{c,K}^{11}$. The lemma is proven.

Lemma 8. *The following equalities hold for the operator $L_{c,K} = L_{c,K}(\lambda)$, $\lambda \in \mathcal{A}_\Theta$:*

$$\begin{aligned} [L_{c,K}]_{21} &= [L_{c-2K,-K}]_{12} \left(\frac{1}{c_2} [L_{c,K}^{11}]_{11} - \frac{c_3}{c_2} \right), \\ [L_{c,K}]_{22} &= [L_{c-2K,-K}]_{11} + \frac{1}{c_2} [L_{c-2K,-K}]_{12} H_{c,K}^{11}. \end{aligned}$$

PROOF. Replace K with $-K$ and c with $c-2K$ in the equality $[L_{c,K}]_{11}\psi_1 + [L_{c,K}]_{12}\psi_2 = \lambda(z)\psi_1$ and multiply both sides by $\frac{\theta(z+2K)}{\theta(z)}$. Observe that d goes into $-d$ and b remains the same upon this change. We obtain

$$[L_{c-2K,-K}]_{11}\psi_2 + [L_{c-2K,-K}]_{12} \frac{\theta(z+2K)\theta(z-2K)}{\theta^2(z)}\psi_1 = \lambda(z)\psi_2.$$

Consequently,

$$\begin{aligned} &[L_{c-2K,-K}]_{12} \left(\frac{1}{c_2} [L_{c,K}^{11}]_{11} - \frac{c_3}{c_2} \right) \psi_1 \\ &+ \left([L_{c-2K,-K}]_{11} + \frac{1}{c_2} [L_{c-2K,-K}]_{12} H_{c,K}^{11} \right) \psi_2 = \lambda(z)\psi_2. \end{aligned}$$

The lemma is proven.

In particular, it follows from Lemma 8 that

$$\begin{aligned} [L_{c,K}^{ij}]_{21} &= H_{c-2K,-K}^{ij} \left(\frac{1}{c_2} [L_{c,K}^{11}]_{11} - \frac{c_3}{c_2} \right), \\ [L_{c,K}^{ij}]_{22} &= [L_{c-2K,-K}^{ij}]_{11} + \frac{1}{c_2} H_{c,K}^{11} H_{c-2K,-K}^{ij}. \end{aligned}$$

Denote the function $H_{c-2K,-K}^{11}$ by V and denote $H_{c,K}^{11}$ by W :

$$V = \frac{2\theta_{11}(K)\theta(c-3K+x)}{\theta(3K)\theta(c-K+x)} \exp \left(-x_1 \left(\frac{b\theta_{11}(K)}{\theta_2(K)} + d \right) \right), \quad (8)$$

$$W = \frac{2\theta_{11}(K)\theta(K+c+x)}{\theta(3K)\theta(c-K+x)} \exp \left(x_1 \left(\frac{b\theta_{11}(K)}{\theta_2(K)} + d \right) \right). \quad (9)$$

From Lemma 7 we obtain

$$h_{c,K}^{11} = \partial_{x_1}^2 \log V + \left(\partial_{x_1} \log V + \frac{3b\theta_{11}(K)}{2\theta_2(K)} + \frac{3d}{2} + \partial_{z_1} \log \theta(3K) \right)^2$$

$$-\frac{\theta_{11}(K)}{\theta_2(K)}(\partial_{x_2} \log V + \partial_{z_2} \log \theta(3K)) + c_3, \quad (10)$$

$$\begin{aligned} h_{c-2K,-K}^{11} &= \partial_{x_1}^2 \log W + \left(\partial_{x_1} \log W - \frac{3b\theta_{11}(K)}{2\theta_2(K)} - \frac{3d}{2} - \partial_{z_1} \log \theta(3K) \right)^2 \\ &\quad + \frac{\theta_{11}(K)}{\theta_2(K)}(\partial_{x_2} \log W - \partial_{z_2} \log \theta(3K)) + c_3. \end{aligned} \quad (11)$$

In a neighborhood of $z = -K$ on Γ_1 we have the expansion

$$\frac{\theta_{12}(z)}{\theta(z)} = \frac{a_2}{\theta_1^2(z)} + \frac{a_1}{\theta_1(z)} + \dots, \quad a_1, a_2 \in \mathbb{C}, \quad (12)$$

and in a neighborhood of $z = K$ on Γ_2 , the expansion

$$\frac{\theta_{12}(z)}{\theta(z)} = \frac{\tilde{a}_2}{\theta_1^2(z)} + \frac{\tilde{a}_1}{\theta_1(z)} + \dots, \quad \tilde{a}_1, \tilde{a}_2 \in \mathbb{C}.$$

As in Lemma 6, we easily demonstrate that $a_1 = -\tilde{a}_1$. Put $a = a_1 = -\tilde{a}_1$.

Lemma 9. *The following hold:*

$$\begin{aligned} h_{c,K}^{12} &= \partial_{x_1} \partial_{x_2} \log V + \left(\partial_{x_1} \log V + \frac{3b\theta_{11}(K)}{2\theta_2(K)} + \frac{3d}{2} + \partial_{z_1} \log \theta(3K) \right) \\ &\quad \times (\partial_{x_2} \log V + \partial_{z_2} \log \theta(3K)) - f_{c,K}^{12} \left(\partial_{x_1} \log V + \frac{3b\theta_{11}(K)}{2\theta_2(K)} + \frac{3d}{2} + \partial_{z_1} \log \theta(3K) \right) \\ &\quad - g_{c,K}^{12} (\partial_{x_2} \log V + \partial_{z_2} \log \theta(3K)) + \partial_{z_1} \partial_{z_2} \log \theta(3K), \\ f_{c,K}^{12} &= \frac{\theta_2(K)}{2\theta_{11}(K)} \left(h_{c,K}^{11} - c_3 + 2 \frac{d\theta_{12}(K)}{\theta_2(K)} - \frac{2b\theta_{11}(K)\theta_{12}(K)}{\theta_2^2(K)} - \frac{2a}{\theta_2(K)} \right. \\ &\quad \left. - \left(\frac{d}{2} - \frac{b\theta_{11}(K)}{2\theta_2(K)} \right)^2 - 2\theta_{11}(K)e - \frac{\theta_{11}(K)}{\theta_2(K)}\alpha \right). \end{aligned}$$

PROOF. In the proof of the lemma we suppose that $z \in \Gamma_1$. Then

$$\begin{aligned} h_{c,K}^{12} &= \partial_{x_1} \partial_{x_2} \log \psi_1 + \partial_{x_1} \log \psi_1 \partial_{x_2} \log \psi_1 - f_{c,K}^{12} \partial_{x_1} \log \psi_1 \\ &\quad - g_{c,K}^{12} \partial_{x_2} \log \psi_1 + \partial_{z_1} \partial_{z_2} \log \theta(z). \end{aligned}$$

Putting $z = -3K$, we hence obtain $h_{c,K}^{12}$.

For convenience, we denote the local parameter $\theta_1(z)$ on Γ_1 at $z = -K$ by k^{-1} . From (3) and (4) we derive the expansion

$$\frac{\theta_1(z)}{\theta(z)} = \frac{\theta_{11}(z)}{\theta_1(z)} - c_3 \frac{\theta(z)}{\theta_1(z)} = \theta_{11}(K)k + d + \frac{e}{k} + o(k^{-1}), \quad e \in \mathbb{C}. \quad (13)$$

Let

$$\frac{\theta_2(z)}{\theta(z)} = \gamma k^2 + \beta k + \alpha + o(1), \quad \alpha, \beta, \gamma \in \mathbb{C}. \quad (14)$$

Then ψ_1 has the form

$$\begin{aligned} \psi_1 &= \frac{1}{\theta(z)} \left(1 + \frac{\xi_1(x)}{k} + \frac{\xi_2(x)}{k^2} + o\left(\frac{1}{k^2}\right) \right) \exp, \\ \exp &= \exp \left(-x_1 \left(\left(\theta_{11}(K)k + d + \frac{e}{k} + \dots \right) - \frac{b\theta_{11}(K)}{2\theta_2(K)} - \frac{d}{2} \right) \right. \\ &\quad \left. - x_2(\gamma k^2 + \beta k + \alpha + \dots) \right). \end{aligned}$$

Equating the coefficients of $k^2 \exp$, $k \exp$, and $\exp$ in (7) to zero, we find that

$$\gamma = -\theta_{11}(K)\theta_2(K), \quad \beta = b\theta_{11}(K) - d\theta_2(K)$$

and

$$h_{c,K}^{11} - c_3 = -2\theta_{11}(K)\partial_{x_1}\xi_1 + \left(\frac{d}{2} - \frac{b\theta_{11}(K)}{2\theta_2(K)} \right)^2 + 2\theta_{11}(K)e + \frac{\theta_{11}(K)}{\theta_2(K)}\alpha.$$

Using (12)–(14), equate the coefficients of $k^2 \exp$ and $k \exp$ in the identity

$$-\partial_{x_1}\partial_{x_2}\psi_1 + f_{c,K}^{12}\partial_{x_1}\psi_1 + g_{c,K}^{12}\partial_{x_2}\psi_1 + h_{c,K}^{12}\psi_1 = \partial_{z_1}\partial_{z_2}\log\theta(z)\psi_1.$$

We obtain $a_2 = \theta_{11}(K)\theta_2(K)$ and

$$f_{c,K}^{12} = -\theta_2(K)\partial_{x_1}\xi_1 + \frac{d\theta_{12}(K)}{\theta_{11}(K)} - \frac{b\theta_{12}(K)}{\theta_2(K)} - \frac{a}{\theta_{11}(K)}.$$

The lemma is proven.

Lemma 10. *The coefficients of the 11-components of the operators $L_{c,K}(\mathcal{A}_\Theta)$ are rationally expressible in terms of V and its derivatives.*

PROOF. Take $z \in \Gamma_1$. Then $[L_{c,K}^{12}]_{11}\psi_1 = \partial_{z_1}\partial_{z_2}\psi_1$. Replace differentiation of ψ_1 with respect to x_2 on the left-hand side of this equality with

differentiation with respect to x_1 (by (7)). We obtain some third-order operator $\tilde{L}$ (in x_1). As follows from Lemmas 7 and 9, the coefficients of $\tilde{L}$ are rationally expressible in terms of V and its derivatives.

Let $[L_{c,K}(\lambda)]_{11}$, $\lambda \in \mathcal{A}_\Theta$, be an arbitrary operator. We have $[L_{c,K}(\lambda)]_{11}\psi_1 = \lambda\psi_1$. As above, replace differentiation with respect to x_2 with differentiation with respect to x_1 . We obtain some operator $\tilde{L}_1$ of order > 3 . The operators $\tilde{L}$ and $\tilde{L}_1$ commute (as having a family of common eigenfunctions parametrized by the points of Γ_1). As demonstrated in [9], the coefficients of $\tilde{L}_1$ are consequently polynomially expressible in terms of the coefficients of $\tilde{L}$ and their derivatives. Hence, the coefficients of $[L_{c,K}(\lambda)]_{11}$ are rationally expressible in terms of V and its derivatives. The lemma is proven.

Similarly, we can prove that the coefficients of the operators $[L_{c-2K,-K}]_{11}$ are rationally expressible in terms of W and its derivatives.

Lemmas 7, 8, and 10 yield the following

Lemma 11. *The coefficients of the second-order Nakayashiki operators are rationally expressible in terms of W , V , and their derivatives.*

The operator $[L_{c,K}^{ijk}]_{11}$ has third order and the principal part $\partial_{x_i}\partial_{x_j}\partial_{x_k}$, whereas the operator $[L_{c,K}^{ijk}]_{12}$ is of the first order [3].

Lemma 12. *The coefficients of the Nakayashiki operators of the third order are rationally expressible in terms of W , V , and their derivatives.*

PROOF. Let $[L_{c,K}]_{12} = u_{c,K}^1\partial_{x_1} + u_{c,K}^2\partial_{x_2} + u_{c,K}^3$, where $L_{c,K}$ is a third-order operator. By Lemma 8, we have

$$\begin{aligned} [L_{c,K}]_{21} &= (u_{c-2K,-K}^1\partial_{x_1} + u_{c-2K,-K}^2\partial_{x_2} + u_{c-2K,-K}^3) \\ &\quad \times \left(\frac{1}{c_2} \left(-\partial_{x_1}^2 + \frac{\theta_{11}(K)}{\theta_2(K)}\partial_{x_2} + h^{11} - c_3 \right) \right), \\ [L_{c,K}]_{22} &= [L_{c-2K,-K}]_{11} + \frac{1}{c_2}(u_{c-2K,-K}^1\partial_{x_1} + u_{c-2K,-K}^2\partial_{x_2} + u_{c-2K,-K}^3)W. \end{aligned}$$

By Lemma 10, the coefficients of $[L_{c,K}]_{11}$ and $[L_{c-2K,-K}]_{11}$ are rationally expressible in terms of W , V , and their derivatives; hence, the coefficient of $\partial_{x_2}^3$ in the 21-component of the commutator $[L_{c,K}^{11}, L_{c,K}] = 0$ has the form

$$-\frac{2\theta_{11}^2(K)}{c_2\theta_2^2(K)}u_{c-2K,-K}^2 + F^2 = 0,$$

where the function F^2 is rationally expressible in terms of W , V , and their derivatives. The coefficients of $\partial_{x_1}\partial_{x_2}^2$ and $\partial_{x_2}^2$ in this component have the

form

$$-\frac{2\theta_{11}^2(K)}{c_2\theta_2^2(K)}u_{c-2K,-K}^i + F^i = 0, \quad i = 1, 3,$$

where the functions F^i are rationally expressible in terms of V , W , $u_{c-2K,-K}^2$, and their derivatives. Consequently, $u_{c-2K,-K}^i$ and $u_{c,K}^i$ are rationally expressible in terms of V , W , and their derivatives. The lemma is proven.

The second- and third-order operators generate the whole ring $L_{c,K}(\mathcal{A}_\Theta)$ [3]; thereby Lemmas 11 and 12 yield Theorem 3.

Write down formulas for the constants c_j (from the introduction).

Multiply both sides of (3) by $\theta^2(z)$ and differentiate with respect to z_2 . Putting $z = K$, we obtain c_2 . Putting $z = 3K$ in (3), we find c_3 . From Lemmas 7 and 9 and formulas (10) and (11) we find the following formulas for the other constants c_j :

$$c_1 = \frac{\theta_{11}(K)}{\theta_2(K)}, \quad c_2 = -\frac{\theta_{11}(K)}{\theta(3K)}, \quad c_3 = \frac{\theta_{11}(3K)}{\theta(3K)} - \frac{\theta_1^2(3K)}{\theta^2(3K)}, \quad (15)$$

$$c_4 = \frac{3}{2}bc_1 + \frac{3d}{2} + \partial_{z_1} \log \theta(3K), \quad c_5 = \partial_{z_2} \log \theta(3K), \quad (16)$$

$$c_6 = \frac{1}{2}bc_1 + \frac{d}{2} + \frac{\theta_{12}(K)}{\theta_2(K)}, \quad c_7 = -c_5c_6 + \partial_{z_1} \partial_{z_2} \log \theta(3K), \quad (17)$$

$$\begin{aligned} c_8 = & 2\frac{d\theta_{12}(K)}{\theta_2(K)} - \frac{2b\theta_{11}(K)\theta_{12}(K)}{\theta_2^2(K)} - \frac{2a}{\theta_2(K)} \\ & - \left(\frac{d}{2} - \frac{b\theta_{11}(K)}{2\theta_2(K)} \right)^2 - 2\theta_{11}(K)e - \frac{\theta_{11}(K)}{\theta_2(K)}\alpha - c_3. \end{aligned} \quad (18)$$

The constants a , b , d , e , and α are determined from (12), (4), (13), and (14).

References.

- [1] Nakayashiki A. Structure of Baker–Akhiezer modules of principally polarized abelian varieties, commuting partial differential operators and associated integrable systems. *Duke Math. J.*, vol. 62, N. 2, 315–358 (1991).
- [2] Nakayashiki A. Commuting partial differential operators and vector bundles over abelian varieties. *Amer. J. Math.*, vol. 116, 65–100 (1994).
- [3] Mironov A. E. Commutative rings of differential operators connected with two-dimensional abelian varieties. *Sibirsk. Mat. Zh.*, vol. 41, N. 6, 1389–1403 (2000).

- [4] Feldman J., Knorrer H., and Trubowitz E. There is no two-dimensional analogue of Lamé's equation. *Math. Ann.*, vol. 294, N. 2, 295–324 (1992).
- [5] Dubrovin B. A. and Natanzon S. M. Real two-gap solutions of the sine-Gordon equation. *Funktsional. Anal. i Prilozhen.*, vol. 16, N. 1, 27–43 (1982).
- [6] Dubrovin B. A. and Novikov S. P. Ground states in a periodic field. Magnetic Bloch functions and vector bundles. *Dokl. Akad. Nauk SSSR*, vol. 253, N. 6, 1293–1297 (1980).
- [7] Mumford D. *Tata Lectures on Theta, Tata Lectures on Theta I, II*, Birkhäuser, Boston-Basel-Stuttgart, 1983, 1984.
- [8] Griffiths F. and Harris J., *Principles of Algebraic Geometry*, N.Y. - Chichester - Brisbane - Toronto : John Wiley & Sons, 1978.
- [9] Krichever I. M. The methods of algebraical geometry in the theory of nonlinear equations. *Uspekhi Mat. Nauk*, vol. 32, N. 6, 183–208 (1997)
- [10] Fay J. D., *Theta functions on Riemann surfaces. Lecture Notes in Math.*; **352**, Springer-Verlag, Berlin, Heidelberg, and New York (1973).
- [11] Taĭmanov I. A. Secants of abelian varieties, theta-functions, and soliton equations. *Uspekhi Mat. Nauk*. vol. 52, N. 1, 149–224 (1997).

Sobolev Institute of Mathematics, Novosibirsk
 mironov@math.nsc.ru